arXiv:1603.09513v1 [math.CA] 31 Mar 2016

# UNCERTAINTY PRINCIPLES FOR THE CLIFFORD-FOURIER TRANSFORM

JAMEL EL KAMEL AND RIM JDAY

Abstract. In this paper, we estabish an analogue of Hardy's theorem and Miyachi's theorem for the Clifford-Fourier transform.

*keywords:* Clifford algebra, Clifford-Fourier transform, Heat kernel, Uncertainty principle, Hardy's theorem, Miyachi's theorem.

MSC (2010) 42B10, 30G35.

## 1. Introduction

Initially, uncertainty principles were given by two theorems: Heisenberg's inequality [19] and Hardy's theorem which ensured that is impossible for a nonzero function and its Fourier transform to be simultaneously small. Afterwards, many theorems were devoted to describe this smallness for example Miyachi [15], Cowling and Price [5], Donoho and Stark [9], Benedicks-Amrein-Berthier [1] and Beurling [2].
Hardy's theorem [11] for the classical Fourier transform states : If we suppose $p$ and $q$ be positive constants and $f$ be a measurable function on the real line satisfying $|f(x)| \leq Ce^{-px^2}$ and $|\mathcal{F}(f)(\lambda)| \leq Ce^{-q\lambda^2}$ for some positive constant C, then (i) $f = 0$ if $pq > \frac{1}{4}$; (ii) $f = Ae^{-px^2}$ for some constant $A$ if $pq = \frac{1}{4}$; (iii) there are many $f$ if $pq < \frac{1}{4}$.
Miyachi's theorem [15] asserts that if $f$ is a measurable function on $\mathbb{R}$ satisfying :

$$e^{ax^2} f \in L^1(\mathbb{R}) + L^\infty(\mathbb{R})$$

and

$$\int_{\mathbb{R}} log^+ \frac{|e^{\frac{y^2}{4a}} \hat{f}(y)|}{\lambda} dy < \infty,$$

for some positive constants $a$ and $\lambda$, then $f$ is a constant multiple of $e^{-ax^2}$.

In Dunkl analysis, analogues of Heisenberg's inequality [16, 17], Hardy's [17], Miyachi's [4], Cowling and Price's [14], Donoho and Stark's [14] and Beurling's [14] theorems were established.
The last decades, there has been an increasing interest in the study of the Clifford-Fourier transform on $\mathbb{R}^m$ introduced in [3] and studied in [7, 8].
In 2010, H. De Bie and Y. Xu in [8] wrote the Clifford-Fourier transform as an integral transform

$$\mathcal{F}_{\pm}(f)(y) = (2\pi)^{\frac{-m}{2}} \int_{\mathbb{R}^m} K_{\pm}(x,y) f(x) dx,$$

where the kernel function $K_{\pm}(x,y)$ was given by an explicit expression. Recently, H. De Bie and Y. Xu in [7] show that the Clifford-Fourier transform is a continuous operator on Schwartz class functions. Besides, they define a translation operator related to the Clifford-Fourier transform and introduce a convolution structure based on translation operator. In the even dimension case, they give an inversion formula for the Clifford-Fourier transform.
Heisenberg's inequality [6, 10], Donoho and Stark's theorem and Benedicks's theorem [13] are obtained for the Clifford-Fourier transform.
The purpose of this paper is to generalize Hardy's theorem and Miyachi's theorem for the Clifford-Fourier transform on $\mathbb{R}^m$.
Our paper is organized as follows. In section 2, we present basic notions and notations related to the Clifford algebra. In section 3, we recall some results for the Clifford-Fourier transform that will be useful in the sequel. Also, we establish some new properties associated to the kernel of the Clifford-Fourier transform as well as for the Clifford-Fourier transform. In section 4, Clifford-heat kernel is introduced and studied. In section 5, we provide Hardy's theorem for the Clifford-Fourier transform on $\mathbb{R}^m$ when $m$ even. Section 6 is devoted to Miyachi's theorem for the Clifford-Fourier transform when $m$ even.
Throughout this paper, the letter C indicates a positive constant that is not necessarily the same in each occurrence.

## 2. Notations and preliminaries

The Clifford algebra $Cl_{0,m}$ over $\mathbb{R}^m$ is a non commutative algebra generated by the basis $\{e_1, .., e_m\}$ satisfying the rules :

$$\begin{cases} e_i e_j = -e_j e_i, & \text{if} \quad i \neq j; \\ e_i^2 = -1, & \forall 1 \leq i \leq m. \end{cases} \tag{2.1}$$

This algebra can be decomposed as

$$Cl_{0,m} = \oplus_{k=0}^{m} Cl_{0,m}^k, \tag{2.2}$$

with $Cl_{0,m}^k$ the space of vectors defined by

$$Cl_{0,m}^k = span\{e_{i_1}..e_{i_k}, i_1 < .. < i_k\}. \tag{2.3}$$

Hence, $\{1, e_1, e_2, .., e_{12}, .., e_{12..m}\}$ forms a basis of $Cl_{0,m}$.
A Clifford number $x$ in $Cl_{0,m}$ is written as follows :

$$x = \sum_{A \in J} e_A x_A, \tag{2.4}$$

where $J := \{0, 1, .., m, 12, .., 12..m\}$, $x_A$ is a real number and $e_A$ belongs to the basis of $Cl_{0,m}$ defined above.
The norm of such element $x$ is given by :

$$||x||_c = \left( \sum_{A \in J} x_A^2 \right)^{\frac{1}{2}}. \tag{2.5}$$

In particular, if $x$ is a vector in $Cl_{0,m}$, then

$$||x||_c^2 = -x^2. \tag{2.6}$$

The Clifford-Dirac operator and Clifford-Laplace operator are defined respectively by :

$$\partial_x = \sum_{i=1}^{m} e_i \partial_{x_i}, \tag{2.7}$$

and

$$\Delta_x = \sum_{i=1}^{m} \partial_{x_i}^2. \tag{2.8}$$

We have the following relation :

$$\Delta_x = -\partial_x^2. \tag{2.9}$$

We introduce respectively the Clifford-Gamma operator associated to a vector $x$, the inner product and the wedge product of two vectors $x$ and $y$ :

$$\Gamma_x := -\sum_{j<k} e_j e_k (x_j \partial_{x_k} - x_k \partial_{x_j}); \tag{2.10}$$

$$< x, y >:= \sum_{j=1}^{m} x_j y_j = -\frac{1}{2}(xy + yx); \tag{2.11}$$

$$x \wedge y := \sum_{j<k} e_j e_k (x_j y_k - x_k y_j) = \frac{1}{2}(xy - yx). \tag{2.12}$$

In the sequel, we consider functions defined on $\mathbb{R}^m$ and taking values in $Cl_{0,m}$. Such functions can be decomposed as :

$$f(x) = f_0(x) + \sum_{i=1}^{m} e_i f_i(x) + \sum_{i<j} e_i e_j f_{ij}(x) + .. + e_1..e_m f_{1..m}(x), \tag{2.13}$$

with $f_0, f_i, .., f_{1..m}$ all real-valued functions.
Let us recall some functional spaces :
$\bullet S(\mathbb{R}^m) \otimes Cl_{0,m}$ the Schwartz space of infinitely differentiable functions on $\mathbb{R}^m$ and taking values in $Cl_{0,m}$ which are rapidly decreasing as their derivatives,
$\bullet \mathcal{P}_k$ the space of homogenious polynomials of degree $k$ taking values in $Cl_{0,m}$,
$\bullet \mathcal{M}_k := \ker \partial_x \cap \mathcal{P}_k$ the space of spherical monogenics of degree $k$,

$\bullet L^p(\mathbb{R}^m)\otimes Cl_{0,m}$ the space of integrable functions taking values in $Cl_{0,m}$ equipped with the norm

$$||f||_{p,c}=\left(\int_{\mathbb{R}^m}||f(x)||_c^p dx\right)^{\frac{1}{p}}=\left(\int_{\mathbb{R}^m}\left(\sum_{A\in J}(f_A(x))^2\right)^{\frac{p}{2}}dx\right)^{\frac{1}{p}}, \tag{2.14}$$

where $J=\{0,1,..,m,12,13,23..,12..m\}$,
$\bullet B(\mathbb{R}^m)\otimes Cl_{0,m}$ a class of integrable functions taking values in $Cl_{0,m}$ and satisfying

$$||f||_{B,c}:=\int_{\mathbb{R}^m}(1+||y||_c)^{\frac{m-2}{2}}||f(y)||_c dy<\infty. \tag{2.15}$$

$\bullet L^\infty(\mathbb{R}^m)\otimes Cl_{0,m}$ the space of essentially bounded functions on $\mathbb{R}^m$ taking values in $Cl_{0,m}$ endowed with the norm

$$||f||_{\infty,c}:=\inf\{C\geq 0;\quad ||f(x)||_c\leq C\quad \text{for almost every } x\in\mathbb{R}^m\}. \tag{2.16}$$

## 3. Clifford-Fourier Transform

**Definition 3.1.** *Let $f\in B(\mathbb{R}^m)\otimes Cl_{0,m}$. The Clifford-Fourier transform is given by* (see [8]) *:*

$$\mathcal{F}_\pm(f)(y)=(2\pi)^{\frac{-m}{2}}\int_{\mathbb{R}^m}K_\pm(x,y)f(x)dx, \tag{3.1}$$

*where*

$$K_\pm(x,y)=e^{\mp i\frac{\pi}{2}\Gamma_y}e^{-i<x,y>}. \tag{3.2}$$

**Lemma 3.1.** *Let $m$ be even. Then*

$$||K_\pm(x,y)||_c\leq Ce^{||x||_c||y||_c},\qquad \forall x,y\in\mathbb{R}^m, \tag{3.3}$$

*with $C$ a positive constant.*

*Proof.* By proposition 3.4 of [8], it is enough to prove the lemma for $K_-(x,y)$.
For $m=2$, it is shown in [8, p.15], that for $x=x_1e_1+x_2e_2$ and $y=y_1e_1+y_2e_2$, we have

$$K_-(x,y)=cos(x_1y_2-x_2y_1)+e_{12}sin(x_1y_2-x_2y_1).$$

Thus

$$||K_-(x,y)||_c=1.$$

For $m>2$. Recall that the kernel $K_-(x,y)$, for $x,y\in\mathbb{R}^m$, can be decomposed as the following

$$K_-(x,y)=K_0^-(x,y)+\sum_{i<j}e_{ij}K_{ij}^-(x,y),$$

with $K_0^-(x,y)$ and $K_{ij}^-(x,y)$ scalar functions,
By lemma 5.2 in [8], for $x,y\in\mathbb{R}^m$,

$$|K_0^-(x,y)|\leq c\,(1+||x||_c||y||_c)^{\frac{m-2}{2}},$$
$$|K_{ij}^-(x,y)|\leq c\,(1+||x||_c||y||_c)^{\frac{m-2}{2}}.$$

Thus

$$|K_-(x,y)| \leq C\,(1+||x||_c||y||_c)^{\frac{m-2}{2}}\,.$$

Since for $u \geq 0$

$$(1+u)^n e^{-u} \leq \frac{n^n}{e^{n-1}}, \forall n \in \mathbb{N},$$

we conclude.

■

**Lemma 3.2.** (see [6] ) *Let $c>0$ and $f \in B(\mathbb{R}^m)\otimes Cl_{0,m}$, then*

$$K_\pm(x,cy) = K_\pm(cx,y), \quad \forall x,y \in \mathbb{R}^m.$$

*Assume that $f_c(x) := f(cx)$, $x \in \mathbb{R}^m$. Then*

$$\mathcal{F}_\pm(f_c)(\lambda) = c^m \mathcal{F}_\pm(f)(c^{-1}\lambda). \tag{3.4}$$

**Theorem 3.3.** (see [7])
*i) The Clifford-Fourier transform defines a continuous operator mapping $S(\mathbb{R}^m)\otimes Cl_{0,m}$ to $S(\mathbb{R}^m)\otimes Cl_{0,m}$* (see [7, Theorem 6.3]).
In particular, when $m$ is even, we have

$$\mathcal{F}_\pm\mathcal{F}_\pm = id_{S(\mathbb{R}^m)\otimes Cl_{0,m}}.$$

ii) The Clifford-Fourier transform extends from $S(\mathbb{R}^m)\otimes Cl_{0,m}$ to a continuous map on $L^2(\mathbb{R}^m)\otimes Cl_{0,m}$.
In particular, when $m$ is even, we have

$$||\mathcal{F}_\pm(f)||_{2,c} = ||f||_{2,c},$$

for all $f \in L^2(\mathbb{R}^m)\otimes Cl_{0,m}$ (see [7, Theorem 6.4]).

**Theorem 3.4.** *Let $\delta>0$ and $P \in \mathcal{P}_k(\mathbb{R}^m)$. Then, there exists $Q \in \mathcal{P}_k(\mathbb{R}^m)$ which satistifies :*

$$\mathcal{F}_\pm(P(.)e^{-\delta||.||_c^2})(x) = Q(x)e^{-\frac{||x||_c^2}{4\delta}}. \tag{3.5}$$

*Proof.* Considering the basis of $L^2(\mathbb{R}^m)\otimes Cl_{0,m}$

$$\phi_{k,l,j}(x) = e^{-\frac{||x||_c^2}{2}} H_{k,m,l}(\sqrt{2}x) P_l^{(j)}(\sqrt{2}x),$$

with $k,l \in \mathbb{N}\cup\{0\}$ and $j=1,..,dim(\mathcal{M}_n^+(l))$.
$H_{k,m,l}$ are the generalized Clifford-Hermite polynomials and

$$\left\{P_l^{(j)}, j=1,2,...,dim(\mathcal{M}_n^+(l)\right\}$$

denotes an orthogonal basis of $\mathcal{M}_n^+(l)$ where $\mathcal{M}_n^+(l)$ the set of all left inner spherical monogenics of degree $l$. This basis constitutes eigenfunction of the Clifford-Fourier transform (see [3]). Note $H_{k,m,l}$ is a polynomial of deree $s$ in the variable $x$ with the real coefficients depending on $k$. Thus, by unicity of eigenfunction of the Clifford-Fourier transform, for every $P \in \mathcal{P}_k(\mathbb{R}^m)$ there exists a unique polynomial $Q$ of degree $k$ such that

$$\mathcal{F}_\pm(P(.)e^{-\frac{||.||_c^2}{2}})(x) = Q(x)e^{-\frac{||x||_c^2}{2}},$$

with degree $Q$ is $k$ .
The proof is completed by lemma 3.2. ■

## 4. Clifford-Heat kernel

In this section, we introduce the Heat kernel in Clifford analysis. Then, we establish some properties of the Clifford-Heat kernel.

**Definition 4.1.** *We define the Clifford-heat kernel by*

$$N_c(x,s) := \frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}}} e^{-\frac{||x||_c^2}{4s}}, \qquad \forall x \in \mathbb{R}^m, s > 0, \tag{4.1}$$

*associated with the Clifford-Laplace operator $\Delta_x$.*

**Theorem 4.1.** *Let $x \in \mathbb{R}^m$ and $s > 0$. Then $N_c(x,s)$ satisfies :*

$$\frac{\partial N_c(x,s)}{\partial s} - \Delta_x N_c(x,s) = 0. \tag{4.2}$$

*Proof.* On one hand, applying the Clifford-Dirac operator to the Clifford-Heat kernel, we obtain

$$\begin{aligned}
\partial_x N_c(x,s) &= \frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}}} \partial_x e^{-\frac{||x||_c^2}{4s}} \\
&= -\frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}+1}} \sum_{i=1}^{m} e_i x_i e^{-\frac{||x||_c^2}{4s}} \\
&= -\frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}+1}} x e^{-\frac{||x||_c^2}{4s}}.
\end{aligned}$$

Thus

$$\begin{aligned}
\partial_x^2 N_c(x,s) &= -\frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}+1}} \partial_x (x e^{-\frac{||x||_c^2}{4s}}) \\
&= -\frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}+1}} (-m - \frac{x^2}{2s}) e^{-\frac{||x||_c^2}{4s}}.
\end{aligned}$$

By (2.6) and (2.9), we get

$$\Delta_x N_c(x,s) = \frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}+1}} (-m + \frac{||x||_c^2}{2s}) e^{-\frac{||x||_c^2}{4s}}.$$

On the other hand, we have

$$\begin{aligned}
\partial_s N_c(x,s) &= \frac{1}{(2\pi)^{\frac{m}{2}}} \partial_s \left( \frac{1}{(2s)^{\frac{m}{2}}} e^{-\frac{||x||_c^2}{4s}} \right) \\
&= \frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}+1}} (-m + \frac{||x||_c^2}{2s}) e^{-\frac{||x||_c^2}{4s}} \\
&= \Delta_c N_x(x,s).
\end{aligned}$$

■

**Theorem 4.2.** *The Clifford-Heat kernel satisfies the following properties :*
i) *For all* $x \in \mathbb{R}^m$ *and* $s > 0$,

$$\mathcal{F}_{\pm}(N_c(.,s))(x) = \frac{1}{(2\pi)^{\frac{m}{2}}} e^{-s||x||_c^2}. \tag{4.3}$$

ii) *Let* $m$ *be even. For* $x \in \mathbb{R}^m$ *and* $s > 0$,

$$N_c(x,s) = \frac{1}{(2\pi)^{\frac{m}{2}}} \int_{\mathbb{R}^m} K_+(y,x) e^{-s||y||_c^2} dy. \tag{4.4}$$

iii) *For all* $\lambda > 0$, $x \in \mathbb{R}^m$ *and* $s > 0$,

$$N_c(\lambda^{\frac{1}{2}}x, \lambda s) = \lambda^{-\frac{m}{2}} N_c(x,s). \tag{4.5}$$

iv) *For* $s > 0$,

$$||N_c(.,s)||_{1,c} = 1. \tag{4.6}$$

v) *For all* $s, t > 0$ *and* $x \in \mathbb{R}^m$,

$$N_c(.,t) *_{Cl} N_c(.,s)(x) = (2\pi)^{\frac{-m}{2}} N_c(x, s+t), \tag{4.7}$$

*where* $*_{Cl}$ *denotes the Clifford-Fourier convolution* (see [8]).

*Proof.* i) One has

$$\begin{aligned}\mathcal{F}_{\pm}(N_c(.,s))(x) &= \frac{1}{(2\pi)^{\frac{m}{2}}} \int_{\mathbb{R}^m} K_{\pm}(y,x) N_c(y,s) dy \\ &= \frac{1}{(2\pi)^m (2s)^{\frac{m}{2}}} \int_{\mathbb{R}^m} K_{\pm}(y,x) e^{-\frac{||y||_c^2}{4s}} dy.\end{aligned}$$

By a change of variable, we get

$$\mathcal{F}_{\pm}(N_c(.,s))(x) = \frac{1}{(2\pi)^{\frac{m}{2}} (2s)^{\frac{m}{2}}} \int_{\mathbb{R}^m} K_{\pm}(\sqrt{2s}z, x) e^{-\frac{||z||_c^2}{2}} dz.$$

Lemma 3.2 implies that

$$\begin{aligned}\mathcal{F}_{\pm}(N_c(.,s))(x) &= \frac{1}{(2\pi)^{\frac{m}{2}}} \int_{\mathbb{R}^m} K_{\pm}(z, \sqrt{2s}x) e^{-\frac{||z||_c^2}{2}} dz \\ &= \mathcal{F}_{\pm}(\phi)(\sqrt{2s}x),\end{aligned}$$

with $\phi(x) = e^{-\frac{||x||_c^2}{2}}$. Since $\mathcal{F}_{\pm}(\phi)(x) = \phi(x)$ (see [8]), we deduce the result.
ii) Using Theorem 3.3 and (4.3), we obtain

$$\begin{aligned}N_c(x,s) = \mathcal{F}_{\pm} \circ \mathcal{F}_{\pm}(N_c(x,s)) &= \frac{1}{(2\pi)^{\frac{m}{2}}} \int_{\mathbb{R}^m} K_{\pm}(y,x) \mathcal{F}_{\pm}(N_c(y,s)) dy \\ &= \frac{1}{(2\pi)^{\frac{m}{2}}} \int_{\mathbb{R}^m} K_{\pm}(y,x) e^{-s||y||_c^2} dy.\end{aligned}$$

iii) Since

$$N_c(\lambda^{\frac{1}{2}}x, \lambda s) = \frac{1}{(2\pi)^{\frac{m}{2}} (2\lambda s)^{\frac{m}{2}}} e^{-\frac{||\lambda^{\frac{1}{2}}x||_c^2}{4\lambda s}},$$

the result is obvious.
iv) By (2.14)

$$||N_c(.,s)||_{1,c} = \int_{\mathbb{R}^m} ||N_c(x,s)||_c dx$$
$$= \frac{1}{(2\pi)^{\frac{m}{2}}(2s)^{\frac{m}{2}}} \int_{\mathbb{R}^m} e^{-\frac{||x||_c^2}{4s}} dx.$$

A change of variable yields the desired result.
v) Since $N_c(.,s)$ is a radial function, then the Clifford-convolution coincides with the classical convolution (see [8]). Thus

$$N_c(.,t) *_{Cl} N_c(.s)(x) = (2\pi)^{-\frac{3m}{2}} 2^{-m} (st)^{-\frac{m}{2}} \int_{\mathbb{R}^m} e^{-\frac{||x-y||_c^2}{4t}} e^{-\frac{||y||_c^2}{4s}} dy.$$

Note that

$$s||x-y||_c^2 = s||x||_c^2 + s||y||_c^2 - 2s < x, y > .$$

$$N_c(.,t) *_{Cl} N_c(.,s)(x) = (2\pi)^{-\frac{3m}{2}} 2^{-m} (st)^{-\frac{m}{2}} e^{-\frac{||x||_c^2}{4(s+t)}} \int_{\mathbb{R}^m} e^{-\frac{||y\sqrt{s+t} - \frac{s}{\sqrt{s+t}}x||_c^2}{4st}} dy.$$

It follows from a change of variable and (4.6) that

$$N_c(.,t) *_{Cl} N_c(.,s)(x) = (2\pi)^{-\frac{3m}{2}} 2^{-\frac{m}{2}} (2st)^{-\frac{m}{2}} e^{-\frac{||x||_c^2}{4(s+t)}} (s+t)^{-\frac{m}{2}} \int_{\mathbb{R}^m} e^{-\frac{||z||_c^2}{4st}} dz$$
$$= (2\pi)^{-m} e^{-\frac{||x||_c^2}{4(s+t)}} (2(s+t))^{-\frac{m}{2}} \int_{\mathbb{R}^m} N_c(z, st) dz$$
$$= (2\pi)^{\frac{-m}{2}} N_c(x, s+t).$$

■

## 5. Hardy's theorem

In this section, we give a generalization of Hardy's theorem for the Clifford-Fourier transform.

**Theorem 5.1.** *Assume that $m$ is even. Let $p$ and $q$ be positive constants. Suppose $f$ is a measurable function on $\mathbb{R}^m$ such that :*

$$||f(x)||_c \leq Ce^{-p||x||_c^2}, \quad x \in \mathbb{R}^m \tag{5.1}$$

*and*

$$||\mathcal{F}_{\pm}(f)(\lambda)||_c \leq Ce^{-q||\lambda||_c^2}, \quad \lambda \in \mathbb{R}^m, \tag{5.2}$$

*for some positive constant $C$. Then, three cases can occur :*

*i) If $pq > \frac{1}{4}$, then $f = 0$.*
*ii) If $pq = \frac{1}{4}$, then $f(x) = Ae^{-p||x||_c^2}$, where $A$ is a constant.*
*iii) If $pq < \frac{1}{4}$, then there are many functions satisfying the assumptions.*

*Proof.* The basis $\{\psi_{j,k,l}\}$ defined in [8] by :

$$\psi_{2j,k,l}(x) := L_j^{\frac{m}{2}+k-1}(||x||_c^2)M_k^{(l)}e^{-\frac{||x||_c^2}{2}},$$

$$\psi_{2j+1,k,l}(x) := L_j^{\frac{m}{2}+k}(||x||_c^2)xM_k^{(l)}e^{-\frac{||x||_c^2}{2}},$$

where $j,k \in \mathbb{N}$, $M_k^{(l)} \in \mathcal{M}_k$; $\quad l = 1,..,dim\mathcal{M}_k$ gives an infinite number of examples for iii).
It is well known that by scaling (lemme 3.2), we may assume $p = q$ without loss of generality. The proof of i) is a simple deduction of ii).

Assume $p = q = \frac{1}{2}$. Since $f$ satisfying (5.1), then $f \in B(\mathbb{R}^m) \otimes Cl_{0,m}$.
Moreover, we have for $\lambda \in \mathbb{R}^m \otimes \mathbb{C}$

$$\begin{aligned}||\mathcal{F}_{\pm}(f)(\lambda)||_c &\leq (2\pi)^{\frac{-m}{2}} \int_{\mathbb{R}^m} ||K_{\pm}(x,\lambda)||_c||f(x)||_c dx \\ &\leq C(2\pi)^{\frac{-m}{2}} \int_{\mathbb{R}^m} ||K_{\pm}(x,\lambda)||_c e^{-\frac{||x||_c^2}{2}} dx.\end{aligned}$$

Applying Lemma 3.1, we get :

$$\begin{aligned}||\mathcal{F}_{\pm}(f)(\lambda)||_c &\leq C(2\pi)^{\frac{-m}{2}} \int_{\mathbb{R}^m} e^{||x||_c||\lambda||_c - \frac{||x||_c^2}{2}} dx \\ &\leq e^{\frac{||\lambda||_c^2}{2}} C(2\pi)^{\frac{-m}{2}} \int_{\mathbb{R}^m} e^{\frac{-(||x||_c - ||\lambda||_c)^2}{2}} dx.\end{aligned}$$

Thus

$$||\mathcal{F}_{\pm}(f)(\lambda)||_c \leq Ce^{\frac{||\lambda||_c^2}{2}}, \tag{5.3}$$

where $C$ is a positive constant.
As $\mathcal{F}_{\pm}(f)$ is an entire function verifying (5.2) and (5.3), lemma 2.1 in [18] allows to express $\mathcal{F}_{\pm}(f)$ as follows :

$$\mathcal{F}_{\pm}(f)(x) = Ae^{-\frac{||x||_c^2}{2}},$$

with A is a constant.
Since $f$ satisfies (5.1), then $f \in L^2(\mathbb{R}^m) \otimes Cl_{0,m}$. By the inversion formula (Theorem 3.3), the proof is completed. ■

## 6. Miyachi's theorem for the Clifford-Fourier transform

In this section, we provide an analogue of Miyachi's theorem for the Clifford-Fourier transform.

**Lemma 6.1.** *Let $m$ be even. Suppose $f$ is a measurable function on $\mathbb{R}^m$ such that*

$$e^{a||.||_c^2} f \in L^p(\mathbb{R}^m) \otimes Cl_{0,m} + L^q(\mathbb{R}^m) \otimes Cl_{0,m}, \tag{6.1}$$

*for some $a > 0$ and $1 \leq p, q \leq +\infty$.*

*Then $\mathcal{F}_{\pm}(f)$ is well defined. Moreover, there exists $C > 0$ such that,*

$$||\mathcal{F}_{\pm}(f)(z)||_c \leq Ce^{||z||_c^2/4a}, \qquad \forall z \in \mathbb{R}^m \otimes \mathbb{C}. \tag{6.2}$$

*Proof.* By (6.1) there exists $u \in L^p(\mathbb{R}^m) \otimes Cl_{0,m}$ and $v \in L^q(\mathbb{R}^m) \otimes Cl_{0,m}$ sucht that

$$f(x) = e^{-a||x||_c^2} \left(u(x) + v(x)\right).$$

Using Hölder's inequality, one has

$$||f||_{B,c} \leq C(||u||_{p,c} + ||v||_{q,c}).$$

Thus, $f \in B(\mathbb{R}^m) \otimes Cl_{0,m}$.
Lemma 3.1 yields for $z \in \mathbb{R}^m \otimes \mathbb{C}$

$$||K_{\pm}(x,z)||_c \leq e^{||x||_c||z||_c}$$

and

$$\begin{aligned} ||\mathcal{F}_{\pm}(f)(z)||_c &\leq C(2\pi)^{\frac{-m}{2}} \int_{\mathbb{R}^m} e^{||x||_c||z||_c} ||f(x)||_c dx \\ &\leq C(2\pi)^{\frac{-m}{2}} e^{\frac{||z||_c^2}{4a}} \int_{\mathbb{R}^m} e^{-a(||x||_c - \frac{||z||_c}{2a})^2} e^{a||x||_c^2} ||f(x)||_c dx. \end{aligned}$$

Applying Hölder's inequality, we deduce that

$$||\mathcal{F}_{\pm}(f)(z)||_c \leq C(2\pi)^{\frac{-m}{2}} e^{\frac{||z||_c^2}{4a}} (||U||_{p,c} + ||V||_{q,c}).$$

Thus, we are done. ∎

**Lemma 6.2.** (see [4]) *Let $h$ be an entire function on $\mathbb{C}^m$ such that*

$$||h(z)|| \leq Ae^{B||Rez||^2}$$

*and*

$$\int_{\mathbb{R}^m} log^+ ||h(y)|| dy < \infty,$$

*for some positive constants $A$ and $B$ where $log^+ x = log\ x$ if $x > 1$ and $log^+ x = 0$ if $x \leq 1$. Then $h$ is a constant.*

**Theorem 6.3.** *Let $m$ be even. Let $a, b$ and $\lambda$ be positive constants and $p$, $q \in [1, +\infty]$. Suppose that $f$ is a measurable function such that*

$$e^{a||.||_c^2} f \in L^p(\mathbb{R}^m) \otimes Cl_{0,m} + L^q(\mathbb{R}^m) \otimes Cl_{0,m} \tag{6.3}$$

*and*

$$\int_{\mathbb{R}^m} log^+ \frac{||e^{b||y||_c^2} \mathcal{F}_{\pm}(f)(y)||_c}{\lambda} dy < \infty. \tag{6.4}$$

*Then, we have the following results :*

- *If $ab > \frac{1}{4}$, then $f = 0$.*
- *If $ab = \frac{1}{4}$, then $f = CN_c(., b)$ with $|C| \leq \lambda$.*

- *If $ab < \frac{1}{4}$, then all functions $f$ written as $f = P(x)N_c(x,\delta)$ with $P \in \mathcal{P}_k$ and $\delta \in ]b, \frac{1}{4a}[$ satisfy the assumptions of the theorem.*

*Proof.* Put

$$h(z) = e^{\frac{-z^2}{4a}}\mathcal{F}_{\pm}(f)(z), \quad \forall z \in \mathbb{R}^m \otimes \mathbb{C}.$$

On one hand, we have for $z = \epsilon + i\eta \in \mathbb{R}^m \otimes \mathbb{C}$

$$z^2 = \epsilon^2 + i\epsilon\eta + i\eta\epsilon - \eta^2 = -||\epsilon||_c^2 - 2i < \epsilon, \eta > +||\eta||_c^2,$$

and

$$||z||_c^2 = ||\epsilon||_c^2 + ||\eta||_c^2.$$

Lemma 6.1 yields

$$||h(\epsilon + i\eta)||_c \leq Ce^{\frac{||\epsilon||_c^2}{2a}}. \tag{6.5}$$

On the other hand, since $log^+(cd) \leq log^+(c) + d$ for all $c, d > 0$, for $ab > \frac{1}{4}$ it follows that

$$\begin{aligned}\int_{\mathbb{R}^m} log^+||h(y)||_c dy &= \int_{\mathbb{R}^m} log^+||e^{\frac{-y^2}{4a}}\mathcal{F}_{\pm}(f)(y)||_c dy = \int_{\mathbb{R}^m} log^+||e^{\frac{||y||_c^2}{4a}}\mathcal{F}_{\pm}(f)(y)||_c dy \\ &= \int_{\mathbb{R}^m} log^+\left(\frac{||e^{b||y||_c^2}\mathcal{F}_{\pm}(f)(y)||_c}{\lambda}\lambda e^{||y||_c^2(\frac{1}{4a}-b)}\right) dy \\ &\leq \int_{\mathbb{R}^m} log^+\frac{||e^{b||y||_c^2}\mathcal{F}_{\pm}(f)(y)||_c}{\lambda} dy + \int_{\mathbb{R}^m} \lambda e^{||y||_c^2(\frac{1}{4a}-b)} dy < +\infty.\end{aligned}$$

As h is an entire function, applying Lemma 6.2, we get $h$ is a constant.
Thus for $ab > \frac{1}{4}$, we have

$$\mathcal{F}_{\pm}(f)(y) = Ce^{\frac{y^2}{4a}} = Ce^{\frac{-||y||_c^2}{4a}}, \forall y \in \mathbb{R}^m.$$

Subsquently, if $ab > \frac{1}{4}$ refering to (6.4), C must be zero.
It is clear that if $f$ satisfies (6.3), then $f \in L^2(\mathbb{R}^m) \otimes Cl_{0,m}$. Thus, Theorem 3.3 implies that

$$f = 0.$$

If $ab = \frac{1}{4}$, then as in the previous case we have

$$\int_{\mathbb{R}^m} log^+\frac{||h(y)||_c}{\lambda} dy = \int_{\mathbb{R}^m} log^+\frac{||e^{b||y||_c^2}\mathcal{F}_{\pm}(f)(y)||_c}{\lambda} dy < +\infty.$$

Thus, we deduce from (6.5) and Lemma 6.2 that

$$\mathcal{F}_{\pm}(f)(y) = Ce^{-b||y||_c^2}.$$

Under condition (6.4), we should have $|c| \leq \lambda$.
Hence, by Theorem 3.3 and (4.3), it follows that

$$f = CN(., b), \text{with } |c| \leq \lambda.$$

For the last case ($ab < \frac{1}{4}$), suppose that $f = P(x)N_c(x,\delta)$ with $\delta \in ]b, \frac{1}{4a}[$ and $P \in \mathcal{P}_k$. Since $\delta < \frac{1}{4a}$, we get

$$e^{a||x||_c^2} f = e^{a||x||_c^2}\psi(x)e^{-\frac{||x||_c^2}{4\delta}} = \psi(x)e^{||x||_c^2(a-\frac{1}{4\delta})} \in L^p(\mathbb{R}^m)\otimes Cl_{0,m} + L^q(\mathbb{R}^m)\otimes Cl_{0,m},$$

where $\psi(x) = \frac{1}{(2\pi)^{\frac{m}{2}}(2\delta)^{\frac{m}{2}}}P(x)$.
Using Theorem 3.4, we obtain

$$\mathcal{F}_{\pm}(f)(x) = Q(x)e^{-\delta||x||_c^2},$$

where $Q$ is a polynomial with the same degree of $P$. Since $b < \delta$ we find that

$$\int_{\mathbb{R}^m} log^+ \frac{||e^{b||y||_c^2}\mathcal{F}_{\pm}(f)(y)||_c}{\lambda}dy = \int_{\mathbb{R}^m} log^+ \frac{||e^{b||y||_c^2}Q(y)e^{-\delta||y||_c^2}||_c}{\lambda}dy < +\infty.$$

■

**Corollary 6.4.** *Let $f$ be a measurable function on $\mathbb{R}^m$ such that*

$$e^{a||.||_c^2} f \in L^p(\mathbb{R}^m)\otimes Cl_{0,m} + L^q(\mathbb{R}^m)\otimes Cl_{0,m} \tag{6.6}$$

$$\int_{\mathbb{R}^m} ||\mathcal{F}_{\pm}(f)(y)||_c^r e^{rb||y||_c^2}dy < \infty, \tag{6.7}$$

*for some constants $a$, $b > 0$, $1 \le p, q \le +\infty$ and $0 < r \le \infty$.*

(i) *If $ab \ge \frac{1}{4}$ then $f = 0$.*
(ii) *If $ab < \frac{1}{4}$ then all functions $f$ of the form $f(x) = P(x)N_c(x,\delta)$ with $P \in \mathcal{P}_k$ and $\delta \in ]b, \frac{1}{4a}[$ satisfy (6.6) and (6.7).*

## References


[1] M. Benedicks, *On Fourier transforms of function supported on sets of finite Lebesgue measure*, J. Math. Anal. Appl., 106 , 180-183.
[2] A. Beurling, *The collect works of Arne Beurling*, Birkhauser. Boston (1989), 1-2.
[3] F. Brackx, N. de Schepper and F.Sommen, *The Clifford-Fourier transform* , J. Fourier Analysis and Applications (1989), Volume 11, Issue 6.
[4] F. Chouchene, R. Daher, T. Kawazoe and H. Mejjaoli, *Miyachi's theorem for the Dunkl transform*, Integral Transforms and Special Functions Vol. 22, No. 3 (2011), 167173.
[5] M.G. Cowling and J.F. Price, *Generalizations of Heisenberg inequality*, Lecture Notes in Math., 992. Springer Berlin (1983), 443-449.
[6] H. De Bie, R. Oste and J. Van der Jeugt, *Generalized Fourier arising from the enveloping algebras of sl(2) and osp(*1|2) Oxford University Press(2015).
[7] H. De Bie, N. De Schepper, F. Sommen, *The Class of Clifford-Fourier Transforms.* J. Fourier Anal Appl,17 (2011):1198-1231 .
[8] H. De Bie and Y. Xu , *On the Clifford-Fourier transform.* J. Fourier Anal. Appl. 11(2005), 669-681.
[9] D.L. Donoho and P.B. Stark, *Uncertainty principles and signal recovery*, SIAM J. Appl. Math., 49 (1989), 906-931.
[10] J. Elkamel R. Jday *The uncertainty principle in Clifford analysis* arXiv:submit/1281037 [math.CA] (16 Jun 2015).
[11] G.H., Hardy, *A theorem concerning Fourier transform*, J. London Math. Soc. 8 (1933), 227-231.
[12] L. Gallardo and K. Trimèche, *An Lp version of Hardys theorem for the Dunkl transform.*

[13] S. Ghobber, Ph. Jaming, *Uncertainty principles for integral operators* arXiv:1206.1195v1 [math.CA] (6 Jun 2012).
[14] T. Kawazoe and H. Mejjaoli, *Uncertainty principles for the Dunkl transform* Hiroshima Mathematical Journal (2010).
[15] A. Miyachi, *A generalization of theorem of Hardy*, Harmonic Analysis Seminar held at Izunagaoka, Shizuoka-Ken, Japon (1997), 44-51.
[16] M. Rosler, *An uncertainty principle for the Dunkl transform*, Bull. Austral. Math. Soc.59 (1999),353-360.
[17] N. Shimeno, *A Note on the Uncertainty Principle for the Dunkl Transform* J. Math. Univ. Tokyo(2001),33-42.
[18] A. Sitaram and M. Sundari, *An analogue of Hardy's theorem for very rapidly decreasing functions on semi-simple Lie groups*, Pacific J. Math. 177(1997), 187-200.
[19] Weyl, *Gruppentheorie and Quantenmechanik*, S. Hirzel, Leipzig,1928. Revized English edition: The Theory of Groups and Qunantum Mechanics, Methuen, London (1931); reprinted by Dover, New York, 1950.

Jamel El Kamel. Department of Mathematics fsm. Monastir 5000, Tunisia.
*E-mail address*: jamel.elkamel@fsm.rnu.tn

Rim Jday. Department of Mathematics fsm. Monastir 5000, Tunisia.
*E-mail address*: rimjday@live.fr